

SHORT COMMUNICATION article

## Evaluation of the antibacterial efficiency of *Carica papaya* leaves against multidrug-resistant *Escherichia coli*

Miriam Goodness U. Nwaneri \*   and Stanley Chikeluom Maduka  

Department of Pharmaceutical Microbiology and Biotechnology, Nnamdi Azikiwe University, Awka,  
Agulu Campus, Anambra State, 5025, Nigeria

\* Author to whom correspondence should be addressed

Received: 27-04-2026, Accepted: 09-06-2026, Published online: 13-06-2026



Copyright© 2026. This open-access article is distributed under the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### HOW TO CITE THIS

Nwaneri MGU, Maduka SC. Evaluation of the antibacterial efficiency of *Carica papaya* leaves against multidrug-resistant *Escherichia coli*. *Mediterr J Med Res.* 2026; 3(2): 201-207. [Article number: 56]. <https://doi.org/10.5281/zenodo.20680923>

**Keywords:** Antioxidant properties, multidrug resistance, phytochemicals

**Abstract:** Over the years, there has been an increase in multidrug-resistant bacteria such as *Escherichia coli*, which has become a serious public health concern globally. It has become obvious that there would possibly be no chemotherapeutic agent to which microorganisms will ultimately grow resistance. This study investigated the in vitro antibacterial activity and the antioxidant activity of the ethanolic extract of *Carica papaya* leaves against multidrug-resistant *Escherichia coli* isolates. The fresh *Carica Papaya* leaves were collected from the plant and air-dried, then the dried leaves were finely ground and soaked for 72 hrs in ethanol to obtain the crude extract. The extract obtained was tested for antibacterial activity against ten *Escherichia coli* isolates using the agar well diffusion method. The phytochemicals and antioxidant properties were tested as well. The isolates were subjected to antibiotic susceptibility testing, which confirmed them all as multidrug-resistant. The findings exhibited concentration-dependent antibacterial activity of the crude extract against the multidrug-resistant *Escherichia coli* isolates, with minimum inhibitory concentration ranging from 25 to 100 mg/mL for most of the isolates. The crude extract showed the presence of alkaloids, flavonoids, terpenoids, and cardiac glycosides. The antioxidant assay for the concentrations 20, 40, 60, 80, and 100 mg/mL indicated a strong radical scavenging ability with percentage values 59.8, 63.2, 69.2, 70.9, and 71.4%, respectively, which indicates that *Carica papaya* leaves possess significant antioxidant properties.

### Introduction

*Escherichia coli* (*E. coli*), a Gram-negative normal intestinal flora of humans and animals, has arisen as one of the most reported pathogens displaying resistance to multiple antibiotics. Infections caused by multidrug-resistant (MDR) *E. coli* strains, such as urinary tract infections, lungs, wounds, septicemia, bone marrow, brain, blood system, and gastrointestinal diseases, have globally led to high morbidity and mortality rates [1]. This constant rise of resistance has pointedly reduced the efficacy of conventional antibiotics, requiring the quest for other healing agents from natural sources. Interest in plant-derived antimicrobial agents has recently intensified due to their availability, safety, and lower tendency to induce resistance compared to synthetic drugs [2]. *Carica papaya*, commonly known as pawpaw or papaya, is a widely cultivated tropical plant belonging to the family Caricaceae.

Various parts of the plant, such as leaves, seeds, fruits, and latex, have been used in old medicine to treat some sicknesses, including malaria, typhoid fever, hypertension, pyrexia, gonorrhoea, diabetes, syphilis, inflammation, and bacterial infections [3, 4]. This study aims to estimate the *in vitro* antibacterial efficiency of *Carica papaya* leaf extracts against MDR *Escherichia coli*.

## Materials and methods

**Culture Media and Reagents:** MacConkey agar, Nutrient agar, Mueller-Hinton agar (MHA), and Nutrient broth (Titan Biotech, India) were the culture media used. They were prepared according to the instructions of the manufacturers. Simmon's citrate (Oxoid, England), peptone water (Titan Biotech, India), hydrogen peroxide (SKG Pharma. Ltd., Nigeria), crystal violet, Lugol's iodine, safranin (May and Baker Ltd., England), Kovac's reagent (Liofilchems, Italy), ethanol, Acetic acid (BDH Chemical Ltd., England) and dimethylsulfoxide (DMSO) were the reagents used.

**Identification and standardization of test microorganisms:** The test organism used in this work includes multidrug-resistant strains of *E. coli* collected from the Laboratory of Pharmaceutical Microbiology and Biotechnology, Nnamdi Azikiwe University, Awka. Each of the test organisms was streaked on a fresh MacConkey agar plate and incubated at 37 °C for 24 hrs. The isolates were identified using macroscopy, microscopy, and confirmatory biochemical tests (oxidase, catalase, indole, and citrate tests) [5, 6]. The bacterial isolates were sub-cultured onto a nutrient agar plate and incubated at 37 °C for 18 - 24 hrs. The inoculum was standardized by transferring a pure culture of the test organism into 3.0 mL of sterile nutrient broth using a sterile wire loop. The suspension was incubated in an oxygen-rich shaker water bath for 3 hrs at 37°C to allow for the growth of the test organism to a density equivalent to the turbidity of 0.5 McFarland [7].

**Antibiotic susceptibility testing:** The susceptibility tests were performed using the M2A6 disc diffusion method [7] as follows. The standardized inocula were swabbed onto a Mueller-Hinton agar plate, and the discs were placed on the inoculated plates and pressed firmly onto the agar plate for complete contact. The plates were left for 30 min to allow for pre-diffusion of antibiotics into the agar and then incubated at 37°C for 18 - 24 hrs. The susceptibility of each isolate to antibiotics was shown by a clear zone of growth inhibition, and the diameter of the zones of inhibition was then interpreted using a standard chart [8].

**Plant material and processing:** Fresh, mature, and fully expanded leaves of *Carica papaya* (**Figure 1**) were collected from the trees growing at the waterside of Agulu Lake, Anambra State, Nigeria. They were identified and authenticated by a plant taxonomist in the Department of Pharmacognosy and Traditional Medicine, Nnamdi Azikiwe University, Awka, Nigeria.



**Figure 1:** *Carica papaya* leaves

The leaves were air dried under shade for some days until properly dried. The dried leaves were milled, and the weight was checked. The grounded leaves were subjected to extraction using Ethyl acetate solvent according to [6] with slight modifications. 85 g of leaves were soaked in 850ml ethyl acetate in a 1000 ml glass beaker, stirred thoroughly, and allowed to stand for 72 hrs to enable complete extraction. It was carefully decanted and placed in a water bath to evaporate the solvent at a reduced temperature of 40 °C. Stock concentrations of 100 mg/ml of the extract were made by weighing 400 mg of the crude extract and reconstituting it in 4 ml DMSO.

*Preliminary antimicrobial assay:* The antimicrobial assay for the plant extract was carried out using an agar well diffusion assay as described. The standardized organisms were inoculated onto previously sterilized MHA plates. A sterile cork borer was used to make five wells (5 mm in diameter) on each of the plates. Two-fold serial dilution was made from the stock concentrations to get graded concentrations of 100, 50, 25, 12.5, and 6.25 mg/ml of the extract. Aliquots of 70 µl of each extract dilution were applied in each of the wells in the culture plates previously seeded with the test organisms. Ciprofloxacin (5.6 mg/ml) served as the positive control against the test organisms. The cultures were incubated at 37 °C for 24 hrs. The antimicrobial activity of each extract was determined by measuring the zone of inhibition around each well. The experiment was done in triplicate against each organism [9, 10].

*Antioxidant assay of crude extract of Carica papaya:* A 16 g of the dried leaf extract was first dissolved in 2.0 ml of DMSO to obtain a concentrated stock solution. From this stock, serial dilutions were prepared to give different concentrations of 100, 80, 60, 40, and 20 mg/ml using methanol as the diluting solvent. The volumes of methanol used for each dilution were 2.70, 2.76, 2.82, 2.88, and 2.94 ml, respectively. A 0.1 mM 2,2-Diphenyl-1-picrylhydrazyl (DPPH) solution was freshly prepared in methanol and protected from light throughout the procedure. The antioxidant reaction mixture for each concentration was prepared by mixing 0.25 ml of each extract dilution with 2.0 ml of the DPPH solution in a clean cuvette. The negative control consisted of methanol mixed with DPPH, while ascorbic acid was used as the positive control. All mixtures were incubated in the dark at room temperature for 30 min to allow complete reaction between the DPPH radicals and the antioxidant components of the extract. After incubation, the absorbance of each sample and control was measured at 517 nm using a spectrophotometer, with methanol serving as the blank. The percentage inhibition of DPPH radicals by the extract was calculated, and the results were noted [11].

*Phytochemical analysis:* The extract was tested for the presence of various plant constituents like alkaloids, flavonoids, reducing sugars, saponins, proteins, tannins, amino acids, steroids, triterpenoids, and glycosides [12-14].

## Results

*Identification of test organisms:* The results of the colonial, microscopic, and biochemical characteristics confirmed them as *Escherichia coli* isolates (**Table 1**).

*Antibiotic susceptibility profile of Escherichia coli isolates:* The antibiotic susceptibility results, as presented in **Tables 2** and **3**, revealed that all isolates exhibited resistance to multiple classes of antibiotics, indicating multidrug resistance.

*Antimicrobial activities of the Carica papaya crude extract against the test organisms:* The result obtained from **Table 3** showed that the extracts have antibacterial activity against the tested *E. coli* isolates. The highest activity was recorded at 100 mg/ml. Isolates EC2, EC3, EC4, EC5, EC6, and EC9 showed inhibition zones ranging from 4 mm to 5 mm.

**Table 1:** Identification of test organisms

Isolate code	Colony morphology	Gram- character	Microscopy	CAT	IND	OXI	CIT
EC 1	Milky, opaque, convex	Gram negative	Rods	+	+	-	-
EC 2	Milky, opaque, convex	Gram negative	Rods	+	+	-	-
EC 3	Milky, opaque, convex	Gram negative	Rods	+	+	-	-
EC 4	Milky, opaque, convex	Gram negative	Rods	+	+	-	-
EC 5	Milky, opaque, convex	Gram negative	Rods	+	+	-	-
EC 3*	Milky, opaque, convex	Gram negative	Rods	+	+	-	-
EC 7	Milky, opaque, convex	Gram negative	Rods	+	+	-	-
EC 8	Milky, opaque, convex	Gram negative	Rods	+	+	-	-
EC 9	Milky, opaque, convex	Gram negative	Rods	+	+	-	-
EC 10	Milky, opaque, convex	Gram negative	Rods	+	+	-	-

**Key:** CAT: catalase, OXI: oxidase, IND: indole, CIT: citrate, EC: *Escherichia coli* tests: + (positive reaction); - (negative reaction)

**Table 2:** Antibiotic susceptibility profile of *Escherichia coli* isolates

Isolate code	Antibiotics/inhibition zone diameter (mm)											
	ACX	CTX	IMP	NF	AUG	OFX	GN	ZEM	NA	CRO	LBC	CXM
EC 1	0±0	0±0	0±0	0 ± 0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0
EC 2	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0
EC 3	0±0	0±0	0±0	12±0	0±0	0±0	9±0	0±0	0±0	14±0.7	15±0	0±0
EC 4	0±0	0±0	0±0	12±0.7	0±0	0±0	0±0	0±0	10±0	20±1.2	16±0	0±0
EC 5	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0
EC 6	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0
EC 7	0±0	0±0	0±0	14±0	0±0	17±0	15±0	0±0	0±0	14±0	16±0.7	0±0
EC 8	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0
EC 9	0±0	0±0	0±0	0±0	0±0	9±1.2	11±0	0±0	0±0	19±0	20±0	0±0
EC 10	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0

**Key:** ACX: Ampiclox (10 µg), CTX: Cefotaxime (25 µg), IMP: Imipenem/Cilastatin (10/10 µg), NF: Nitrofurantoin (300 µg), AUG: Amoxicillin Clavulanate (30 µg), OFX: Ofloxacin (5 µg), GN: Gentamycin (10 µg), ZEM: Cefexime (5 µg), NA: Nalidixic Acid (30 µg), CRO: Ceftriaxone Sulbactam (45 µg), LBC: Levofloxacin (5 µg); CXM: Cefuroxime (30 µg), EC (*Escherichia coli*); 0±0 no inhibition zone.

**Table 3:** Antimicrobial activities of the *Carica papaya* Crude extract against the test organisms

Conc. (mg/mL)	Test organisms/inhibition zone diameter (mm)									
	EC1	EC2	EC3	EC4	EC5	EC6	EC7	EC8	EC9	EC10
100	0±0	5±0	4±0	5±0	4±0	5±0	0±0	0±0	5±0	3±0
50	0±0	4±0.7	3±0.7	3±0.7	3±0	4±0	0±0	0±0	4±0	0±0
25	0±0	4±0	3±0	3±0	0±0	3±0	0±0	0±0	2.5±0	0±0
12.5	0±0	3±0	1.5±0	2.5±0	0±0	2±0	0±0	0±0	2±0	0±0
6.25	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0	0±0

**Key:** EC (*Escherichia coli*), 0±0 no inhibition zone.

**Antioxidant assay of crude extract of *Carica papaya*:** The crude extract of *Carica Papaya* showed a percentage inhibition which ranged from 59.8% at 20 mg/ml to 71.4% at 100 mg/ml, indicating that antioxidant activity increased with an increase in concentration (**Table 4**).

**Phytochemical assay of *Carica papaya* extract:** The result of the phytochemical screening of the *Carica papaya* leaf extract (**Table 5**) revealed the presence of Alkaloids, flavonoids, terpenoids, and cardiac glycosides, indicating the extract antimicrobial, antioxidant, and therapeutic activities [22]. However, reducing sugars, saponins, tannins, and proteins were not detected in the extract.

**Table 4:** Antioxidant assay of crude extract of *Carica papaya*

Concentration (mg/mL)	<i>Carica papaya</i>	Ascorbic acid
	Mean absorbance / % Inhibition	Mean absorbance / % Inhibition
100	0.230 (71.4)	0.200 (87.5)
80	0.233 (70.9)	0.205 (82.9)
60	0.248 (69.2)	0.211 (77.7)
40	0.296 (63.2)	0.216 (73.6)
20	0.323 (59.8)	0.220 (70.45)

**Table 5:** Phytochemical assay of *Carica papaya* extract

Plant	Alkaloid	Flavonoid	Terpenoid	Reducing sugars	Cardiac glycosides	Saponin	Tanin	Protein
<i>Carica papaya</i>	+	+	+	-	+	-	-	-

## Discussion

The high resistance of the bacterial isolates to antibiotics agrees with reports that *Escherichia coli* rapidly acquires antibiotic resistance due to horizontal gene transfer and antibiotic misuse [15-17]. The antibacterial activity of the extract supports previous findings of [18], which revealed that *Carica papaya* is effective against resistant *E. coli* strains. The inhibition zone diameter decreased as the extract concentration decreased, which indicates a concentration-dependent antibacterial response. Conversely, EC1, EC7, EC8, and EC10 showed no inhibition zone diameter, which indicates resistance. The results confirm that *Carica papaya* possesses notable antioxidant potential and free radical scavenging ability. This can counter pro-oxidants through several signaling pathways that either help the expression of antioxidant enzymes or reduce ROS production. As such, it can be combined into supplements and medications to help handle health situations motivated by oxidative stress. Our work agrees with the studies of [19, 20] that reported the plant's antioxidant properties. The presence of these bioactive compounds gives *Carica papaya* leaves their medicinal value, as alkaloids and flavonoids are known for their strong antimicrobial and antioxidant properties [21], while terpenoids and cardiac glycosides contribute to antibacterial and anti-inflammatory activities [22].

**Conclusion:** This study demonstrates that the ethanolic extract of *Carica papaya* leaves possesses antibacterial and antioxidant effects against several multidrug-resistant *Escherichia coli* isolates within the environment, with activity increasing in a concentration-dependent manner. Their efficiency could be due to the presence of alkaloids, flavonoids, terpenoids, and cardiac glycosides found in the extract. We therefore recommend further studies on the extract to determine its safety and toxicity profile. Also, clinical studies should be conducted to assess its potential use for the management of resistant *Escherichia coli* infections.

## References

1. Shiela C. *Escherichia coli*: An arduous voyage from commensal to antibiotic resistance. *Microbial Pathogenesis*. 2025; 198: 107173. doi: 10.1016/j.micpath.2024.107173
2. Álvarez-Martínez FJ, Barraón-Catalán E, Micol V. Tackling antibiotic resistance with compounds of natural origin: A comprehensive review. *Biomedicines*. 2020; 8(10): 405. doi: 10.3390/biomedicines8100405
3. Arora R, Malhotra P, Sharma A, Haniadka R, Yashawanth HS, Baliga MS. Medicinal efficacy of Indian herbal remedies for the treatment of arthritis. *Bioactive food as dietary interventions for arthritis and related inflammatory diseases*, Academic Press. 2013; 601-617. doi: 10.1016/B978-0-12-397156-2.00250-7
4. Vij T, Prashar Y. A review on medicinal properties of *Carica papaya* Linn. *Asian Pacific Journal of Tropical Disease*. 2015; 5(1): 1-6. doi: 10.1016/S2222-1808(14)60617-4
5. Ekwealor PA, Ugwu MC, Ezeobi I, Amalukwe G, Ugwu BC, Okezie U, et al. Antimicrobial evaluation of bacterial isolates from urine specimens of patients with urinary tract infections in Awka, Nigeria. *International Journal of Microbiology*. 2016; 9740273. doi: 10.1155/2016/9740273
6. Ebenebe IN, Nedum CH, Okezie UM, Egbuna NR, Obasi CC, Nwaneri MGU. Comparative assessment of *Solanum melongena* (Eggplant) against multi-drug-resistant *Staphylococcus aureus* and *Pseudomonas aeruginosa*. *Mediterranean Journal of Pharmacy and Pharmaceutical Sciences*. 2024; 4(4): 33-40. doi: 5281/zenodo.14176439
7. Nwaneri MGU, Ilechukwu OA, Ugo UA. Antimicrobial activity of endophytic fungal metabolites from breadfruit (*Artocarpus altilis*) leaves against multidrug-resistant *Staphylococcus aureus* and *Klebsiella spp.* *Mediterranean Journal of Medicine and Medical Sciences*. 2025; 1(3): 23-30. doi: 10.5281/zenodo.17611720
8. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing; 24<sup>th</sup> informational supplement. Clinical and Laboratory Standards Institute; Wayne, PA: 2014; (CLSI document M100-S24).
9. Okezie UM, Obi MC, Morikwe UC, Ebenebe IN, Nwaneri MGU. Antimicrobial and antioxidant potentials of crude extracts of culturally dissimilar endophytic fungi. *GSC Biological and Pharmaceutical Sciences*. 2023; 22(2): 187-195. doi: 10.30574/gscbps.2023.22.2.0475
10. Nwaneri MGU, Umeogoji LN, Ugo UA. Fungicidal action of endophytic fungi, obtained from *Justicia carnea*, against multidrug-resistant *Candida albicans*. *Mediterranean Journal of Medical Research*. 2025; 2(4): 179-185. doi: 10.5281/zenodo.17449274
11. Ebenebe IN, Uzor KE, Okezie UM, Chukwunwejim CR, Nwaneri MGU, Egbuna NR, Eze PM. Antimicrobial and antioxidant potentials of an endophytic *Cunninghamella* sp. isolated from the leaves of *Chrysophyllum albidum*. *Magna Scientia Advanced Biology and Pharmacy*. 2023; 09(01): 071-076. doi.org/10.30574/msabp.2023.9.1.0036
12. Kokate CK. A textbook of practical pharmacognosy. 5<sup>th</sup> Ed., New Delhi: Vallabh Prakashan, India. 2005. ISBN: 978-9386310002.
13. Harborne JB. Phytochemical methods: A guide to modern techniques of plant analysis. 5<sup>th</sup> Ed., London: Chapman and Hall. 1998. ISBN: 9780412572609.
14. Khandewal KR. Practical pharmacognosy. Nirali Prakashan, Pune; 4<sup>th</sup> Ed., Pragati Books Pvt. Ltd., New Delhi, India. 2008. ISBN: 9788185790305.
15. Kerek Á, Román I, Szabó Á, Kovács D, Kardos G, Kovács L, Jerzsele Á. Antibiotic resistance genes in *Escherichia coli*: A literature review. *Critical Reviews in Microbiology*. 2026; 52(1): 1-35. doi: 10.1080/1040841X.2025.2492156
16. Perry J, Waglechner N, Wright G. The prehistory of antibiotic resistance. *Cold Spring Harb Perspect Med*. 2016; 6(6): a025197. doi: 10.1101/cshperspect.a025197
17. Nasrollahian S, Graham JP, Halaji M. A review of the mechanisms that confer antibiotic resistance in pathotypes of *E. coli*. *Frontiers in Cell. Infect. Microbiology*. 2024; 14: 1387497. doi: 10.3389/fcimb.2024.1387497
18. Alhodieb FS, Farid M, Sabir M, Nisa S, Sarwar S, Abbas S. Exploring the bioactive compounds of *Carica papaya* leaves: phytol's role in combating antibiotic-resistant bacteria. *Frontiers in Cellular and Infection Microbiology*. 2025; 15: 1564787. doi: 10.3389/fcimb.2025.1564787
19. Kong YR, Jong YX, Balakrishnan M, Bok ZK, Weng JKK, Tay KC, et al. Beneficial role of carica papaya extracts and phytochemicals on oxidative stress and related diseases: A mini review. *Biology (Basel)*. 2021; 10(4): 287. doi: 10.3390/biology10040287
20. Sharma AS, Ruchi SM, Kumar MB, Mrunal Deepak, L, José M, et al. *Carica papaya* L. Leaves: Deciphering Its antioxidant bioactives, biological activities, innovative products, and safety aspects. *Oxidative Medicine and Cellular Longevity*. 2022: 2451733. doi: 10.1155/2022/2451733

21. Alara OR, Abdurahman NH, Alara JA. *Carica papaya*: Comprehensive overview of the nutritional values, phytochemicals and pharmacological activities. *Advances in Traditional Medicine*. 2022; 22: 17-47. doi: 10.1007/s13596-020-00481-3
22. Sabir K, Zeemal SA, Bushra SA, Muhammed TM, Katarzyna KB, et al. Phytochemical profiling, antioxidant potential, and in silico evaluation of *Carica papaya* seed oil. *Results in Chemistry*. 2025; 18:102751. doi: 10.1016/j.rechem.2025.102751

**Acknowledgments:** The authors are thankful to the Faculty of Pharmaceutical Sciences, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria.

**Authors' Contributions:** Both authors conceptualized and designed the study; SCM conducted data collection; MGUN conducted data analysis. Both authors are writing, editing, and proofreading. Both authors read and approved the manuscript.

**Conflict of interest:** The authors declare the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Ethical issues:** Both authors are responsible for ethical issues, including plagiarism, informed consent, data fabrication or falsification, and duplicate publication or submission.

**Data availability statement:** The raw data that support the findings of this article are available from the corresponding author upon reasonable request.

**Author declarations:** The authors confirm that they have followed all relevant ethical guidelines and obtained any necessary IRB and/or ethics committee approvals.

**Generative AI disclosure:** No generative AI was used in the preparation of this manuscript.

## تقييم فعالية أوراق البابايا (*Carica papaya*) كمضاد للبكتيريا ضد بكتيريا الإشريكية القولونية المقاومة للأدوية المتعددة

ميريام جودنس يو. نوانيري\* وستانلي تشيكيكوم مادوكا

قسم علم الأحياء الدقيقة الصيدلانية والتكنولوجيا الحيوية، جامعة نامدي أزيكيوي، أوكا،

حرم أغولو، ولاية أنامبرا، 5025، نيجيريا  
\* المؤلف المسؤول عن المراسلات

ملخص: على مر السنين، ازداد انتشار البكتيريا المقاومة للأدوية المتعددة، مثل الإشريكية القولونية، مما أصبح مصدر قلق بالغ للصحة العامة على مستوى العالم. بات من الواضح أنه من غير المرجح وجود أي عامل علاجي كيميائي يمكن للكائنات الدقيقة أن تُطور مقاومة له في نهاية المطاف. هدفت هذه الدراسة إلى التحقق من النشاط المضاد للبكتيريا والنشاط المضاد للأكسدة لمستخلص الإيثانول من أوراق البابايا (*Carica papaya*) ضد عزلات الإشريكية القولونية المقاومة للأدوية المتعددة. جُمعت أوراق البابايا الطازجة من النبات وجُففت هوائياً، ثم طُحنت الأوراق المجففة طحناً ناعماً ونُقتعت لمدة 72 ساعة في الإيثانول للحصول على المستخلص الخام. خضع المستخلص لاختبار النشاط المضاد للبكتيريا ضد عشر عزلات من الإشريكية القولونية باستخدام طريقة الانتشار في الآبار على وسط الآجار. كما تم اختبار المركبات الكيميائية النباتية وخصائص مضادات الأكسدة. خضعت العزلات لاختبار حساسية المضادات الحيوية، والذي أكد مقاومتها جميعاً للأدوية المتعددة. أظهرت النتائج نشاطاً مضاداً للبكتيريا يعتمد على التركيز للمستخلص الخام ضد عزلات الإشريكية القولونية المقاومة للأدوية المتعددة، حيث تراوح الحد الأدنى للتركيز المثبط من 25 إلى 100 ملغم/مل لمعظم العزلات. وكشف المستخلص الخام عن وجود قلويدات، وفلافونويدات، وتربينويدات، وجليكوسيدات قلبية. وأشارت نتائج اختبار مضادات الأكسدة للتركيزات 20، 40، 60، 80، و100 ملغم/مل إلى قدرة قوية على إزالة الجذور الحرة بنسب مئوية بلغت 59.8%، 63.2%، 69.2%، 70.9%، و71.4% على التوالي، مما يدل على أن أوراق البابايا (*Carica papaya*) تمتلك خصائص مضادة للأكسدة فعالة.